Pawan Jaiswal, Vikram Singh and Manoj Kumar Shukla

Received November 25, 2016 and Accepted February 22, 2017

ABSTRACT: The field experiment was conducted at Crop Research Farm, Department of Agronomy, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad, U.P., during *kharif* season of 2011. The experiment consisted of three levels of nitrogen (0 kg/ha, 20 kg/ha and 30 kg/ha) and three levels of phosphorus (30 kg/ha, 45 kg/ha and 60 kg/ha) with *Rhizobium* inoculation and uninoculation. Highest plant height, plant dry weight, Number of nodules, Crop Growth Rate and Relative Growth Rate was found in the treatment (T_{10}) Nitrogen 20 kg/ha + Phosphorus 45 kg/ha in greengram than other treatments. More number of pod/plant, seed/pod, highest test weight, seed yield, stover yield and harvest index was recorded also in above respective treatment. Maximum Gross return, Net return and Benefit cost ratio were recorded in T_{10} (Nitrogen 20 kg/ha + Phosphorus 45 kg/ha + inoculated *Rhizobium*) Rs./ha.

Pawan Jaiswal, Vikram Singh and Manoj Kumar Shukla

Received November 25, 2016 and Accepted February 22, 2017

ABSTRACT: The field experiment was conducted at Crop Research Farm, Department of Agronomy, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad, U.P., during *kharif* season of 2011. The experiment consisted of three levels of nitrogen (0 kg/ha, 20 kg/ha and 30 kg/ha) and three levels of phosphorus (30 kg/ha, 45 kg/ha and 60 kg/ha) with *Rhizobium* inoculation and uninoculation. Highest plant height, plant dry weight, Number of nodules, Crop Growth Rate and Relative Growth Rate was found in the treatment (T_{10}) Nitrogen 20 kg/ha + Phosphorus 45 kg/ha in greengram than other treatments. More number of pod/plant, seed/pod, highest test weight, seed yield, stover yield and harvest index was recorded also in above respective treatment. Maximum Gross return, Net return and Benefit cost ratio were recorded in T_{10} (Nitrogen 20 kg/ha + Phosphorus 45 kg/ha + inoculated *Rhizobium*) Rs./ha.

Pawan Jaiswal, Vikram Singh and Manoj Kumar Shukla

Received November 25, 2016 and Accepted February 22, 2017

ABSTRACT: The field experiment was conducted at Crop Research Farm, Department of Agronomy, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad, U.P., during *kharif* season of 2011. The experiment consisted of three levels of nitrogen (0 kg/ha, 20 kg/ha and 30 kg/ha) and three levels of phosphorus (30 kg/ha, 45 kg/ha and 60 kg/ha) with *Rhizobium* inoculation and uninoculation. Highest plant height, plant dry weight, Number of nodules, Crop Growth Rate and Relative Growth Rate was found in the treatment (T_{10}) Nitrogen 20 kg/ha + Phosphorus 45 kg/ha in greengram than other treatments. More number of pod/plant, seed/pod, highest test weight, seed yield, stover yield and harvest index was recorded also in above respective treatment. Maximum Gross return, Net return and Benefit cost ratio were recorded in T_{10} (Nitrogen 20 kg/ha + Phosphorus 45 kg/ha + inoculated *Rhizobium*) Rs./ha.

Pawan Jaiswal, Vikram Singh and Manoj Kumar Shukla

Received November 25, 2016 and Accepted February 22, 2017

ABSTRACT: The field experiment was conducted at Crop Research Farm, Department of Agronomy, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad, U.P., during *kharif* season of 2011. The experiment consisted of three levels of nitrogen (0 kg/ha, 20 kg/ha and 30 kg/ha) and three levels of phosphorus (30 kg/ha, 45 kg/ha and 60 kg/ha) with *Rhizobium* inoculation and uninoculation. Highest plant height, plant dry weight, Number of nodules, Crop Growth Rate and Relative Growth Rate was found in the treatment (T_{10}) Nitrogen 20 kg/ha + Phosphorus 45 kg/ha in greengram than other treatments. More number of pod/plant, seed/pod, highest test weight, seed yield, stover yield and harvest index was recorded also in above respective treatment. Maximum Gross return, Net return and Benefit cost ratio were recorded in T_{10} (Nitrogen 20 kg/ha + Phosphorus 45 kg/ha + inoculated *Rhizobium*) Rs./ha.

Pawan Jaiswal, Vikram Singh and Manoj Kumar Shukla

Received November 25, 2016 and Accepted February 22, 2017

ABSTRACT: The field experiment was conducted at Crop Research Farm, Department of Agronomy, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad, U.P., during *kharif* season of 2011. The experiment consisted of three levels of nitrogen (0 kg/ha, 20 kg/ha and 30 kg/ha) and three levels of phosphorus (30 kg/ha, 45 kg/ha and 60 kg/ha) with *Rhizobium* inoculation and uninoculation. Highest plant height, plant dry weight, Number of nodules, Crop Growth Rate and Relative Growth Rate was found in the treatment (T_{10}) Nitrogen 20 kg/ha + Phosphorus 45 kg/ha in greengram than other treatments. More number of pod/plant, seed/pod, highest test weight, seed yield, stover yield and harvest index was recorded also in above respective treatment. Maximum Gross return, Net return and Benefit cost ratio were recorded in T_{10} (Nitrogen 20 kg/ha + Phosphorus 45 kg/ha + inoculated *Rhizobium*) Rs./ha.

Pawan Jaiswal, Vikram Singh and Manoj Kumar Shukla

Received November 25, 2016 and Accepted February 22, 2017

ABSTRACT: The field experiment was conducted at Crop Research Farm, Department of Agronomy, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad, U.P., during *kharif* season of 2011. The experiment consisted of three levels of nitrogen (0 kg/ha, 20 kg/ha and 30 kg/ha) and three levels of phosphorus (30 kg/ha, 45 kg/ha and 60 kg/ha) with *Rhizobium* inoculation and uninoculation. Highest plant height, plant dry weight, Number of nodules, Crop Growth Rate and Relative Growth Rate was found in the treatment (T_{10}) Nitrogen 20 kg/ha + Phosphorus 45 kg/ha in greengram than other treatments. More number of pod/plant, seed/pod, highest test weight, seed yield, stover yield and harvest index was recorded also in above respective treatment. Maximum Gross return, Net return and Benefit cost ratio were recorded in T_{10} (Nitrogen 20 kg/ha + Phosphorus 45 kg/ha + inoculated *Rhizobium*) Rs./ha.

Pawan Jaiswal, Vikram Singh and Manoj Kumar Shukla

Received November 25, 2016 and Accepted February 22, 2017

ABSTRACT: The field experiment was conducted at Crop Research Farm, Department of Agronomy, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad, U.P., during *kharif* season of 2011. The experiment consisted of three levels of nitrogen (0 kg/ha, 20 kg/ha and 30 kg/ha) and three levels of phosphorus (30 kg/ha, 45 kg/ha and 60 kg/ha) with *Rhizobium* inoculation and uninoculation. Highest plant height, plant dry weight, Number of nodules, Crop Growth Rate and Relative Growth Rate was found in the treatment (T_{10}) Nitrogen 20 kg/ha + Phosphorus 45 kg/ha in greengram than other treatments. More number of pod/plant, seed/pod, highest test weight, seed yield, stover yield and harvest index was recorded also in above respective treatment. Maximum Gross return, Net return and Benefit cost ratio were recorded in T_{10} (Nitrogen 20 kg/ha + Phosphorus 45 kg/ha + inoculated *Rhizobium*) Rs./ha.